



YOU'RE INVITED:

JOIN US FOR A FREE *Cool* EVENT

DISCOVER HOW COOLSCULPTING® CAN HELP YOU ELIMINATE STUBBORN FAT

8 CoolSculpting treatments on the abdomen over 2 visits



BEFORE

4 WEEKS AFTER 1ST VISIT

8 WEEKS AFTER 2ND VISIT

Before-during-after photos courtesy of Bodify® and Steven Sorr, NMD. Results and patient experience may vary.

Join us for a CoolEvent to learn more about CoolSculpting and if it's the right treatment for you.

HOSTED BY:

ADDRESS:

DATE AND TIME:

SPECIAL OFFER:

RESERVE YOUR SPOT TODAY!

Uses
CoolSculpting® and CoolSculpting® Elite are FDA-cleared for the treatment of visible fat bulges in the submental (under the chin) and submandibular (under the jawline) areas, thigh, abdomen, and flank, along with bra fat, back fat, underneath the buttocks (also known as banana roll), and upper arm. It is also FDA-cleared to affect the appearance of lax tissue with submental area treatments. CoolSculpting® and CoolSculpting® Elite are not treatments for weight loss.

Important Safety Information
These procedures are not for everyone. You should not be treated with CoolSculpting® or CoolSculpting® Elite if you suffer from cryoglobulinemia, cold agglutinin disease, or paroxysmal cold hemoglobinuria.

Tell your doctor if you have any medical conditions including recent surgery, pre-existing hernia, and any known sensitivities or allergies.

During the procedure you may experience sensations of pulling, tugging, mild pinching, intense cold, tingling, stinging, aching, and cramping at the treatment site. These sensations lessen as the area becomes numb. Following the procedure, typical side effects include temporary redness, swelling, blanching, bruising, firmness, tingling, stinging, tenderness, cramping, aching, itching, or skin sensitivity, and sensation of fullness in the back of the throat after submental or submandibular area treatment.

Rare side effects may happen in 1 to 10 out of 10,000 CoolSculpting® and CoolSculpting® Elite treatments (between 0.01% to 0.1%). One such rare side effect is a visible enlargement in the treated area, which may develop 2 to 5 months after treatment, will not resolve on its own, and may require surgical intervention for correction.

Please see full Important Safety Information for CoolSculpting® and CoolSculpting® Elite on CoolSculpting.com.